arXiv:math/0702001v1 [math.QA] 31 Jan 2007

K-THEORETIC CONSTRUCTION OF NONCOMMUTATIVE INSTANTONS OF ALL CHARGES

Ludwik Dąbrowski
Scuola Internazionale Superiore di Studi Avanzati
Via Beirut 2-4, Trieste, 34014 Italy
e-mail: dabrow@sissa.it

Tom Hadfield
School of Mathematical Sciences, Queen Mary, University of London
327 Mile End Road, London E1 4NS, United Kingdom
e-mail: t.hadfield@qmul.ac.uk

Piotr M. Hajac
Instytut Matematyczny, Polska Akademia Nauk
ul. Śniadeckich 8, Warszawa, 00-956 Poland
http://www.impan.gov.pl/~pmh
and
Katedra Metod Matematycznych Fizyki, Uniwersytet Warszawski
ul. Hoża 74, Warszawa, 00-682 Poland

Rainer Matthes
Katedra Metod Matematycznych Fizyki, Uniwersytet Warszawski
ul. Hoża 74, Warszawa, 00-682 Poland
e-mail: matthes@fuw.edu.pl, matthes@itp.uni-leipzig.de

February 2, 2008

Abstract

We construct noncommutative versions of instanton bundles of arbitrary charges.

1 Introduction

The goal of this paper is to construct noncommutative analogues of the classical instanton vector bundles of arbitrary charge [At] associated to the Hopf fibration

$$SU(2) \to S^7 \to S^4 \tag{1.1}$$

In Connes' framework for noncommutative geometry the concept of a noncommutative vector bundle is well understood: by the Serre-Swan theorem, these correspond to finitely-generated projective modules over a C*-algebra representing the underlying noncommutative space. The concept of a noncommutative principal bundle is less well-established, although recently there have been substantial developments of the latter both concerning its clarification and the construction of interesting examples. In this paper we present a construction of finitely-generated projective modules over a C*-algebra corresponding to a noncommutative 4-sphere, which reduce to classical instanton bundles of arbitrary charge in the commutative situation. The problem of giving a corresponding noncommutative analogue of (1.1), in particular finding a noncommutative 7-sphere equipped with an appropriate action of quantum $SU(2)$, will be treated elsewhere.

Recall that classically the Hopf fibration (1.1) with winding number $n \in \mathbb{Z}$ can be described as follows. Consider S^4 as the one-point compactification of $\mathbb{R}^4$, with the open covering

$$U_0 = \{\ \mathbf{x} = (x_0, x_1, x_2, x_3)\ :\ ||\mathbf{x}|| < 1 + \varepsilon\ \}, \quad U_1 = \{\ \mathbf{x}\ :\ ||\mathbf{x}|| > 1 - \varepsilon\ \}$$

for some $\varepsilon > 0$. Then $U_{01} := U_0 \cap U_1 \cong S^3 \times (1-\varepsilon, 1+\varepsilon)$. We identify $S^3 = \{\ \mathbf{x} = (x_0, x_1, x_2, x_3)\ :\ ||\mathbf{x}|| = 1\ \}$ with $SU(2)$ via the map $\mathbf{x} \mapsto x_0 I_2 + \sum_{i=1}^{3} x_i \sigma_i$ where the $\sigma_i \in M_2(\mathbb{C})$ are the standard Pauli matrices. Any transition function $U_{01} \to U_{01}$ defines a continuous map from S^3 to $SU(2)$, hence these are classified by $\pi_3(S^3) \cong \mathbb{Z}$. The transition function $\theta^{(n)}$ corresponding to winding number $n \in \mathbb{Z}$ is

$$\theta^{(n)}(\mathbf{x}) = (x_0 I_2 + \sum_{i=1}^{3} x_i \sigma_i)^n \tag{1.2}$$

Various quantum deformations of (1.1) with structure group $SU(2)$ replaced by Woronowicz's quantum group $SU_q(2)$ [Wo87] exist in the literature. The earliest example known to us is the sheaf-theoretic construction of Pflaum [Pfl], although the coaction of the structure group is not self-evident therein. Two relevant constructions of quantum 4-spheres, given by C*-algebras denoted $C(S^4_\theta)$ and $C(S^4_q)$ were found in [CL01, DLM]. The second of these is a suspension of $SU_q(2)$, viewed as a quantum 3-sphere (the first being a suspension of the Matsumoto quantum 3-sphere [Mat]), and coincides with the quantum 4-sphere B which we use in this paper. Based on this work, quantum versions of (1.1) were given in [LPR] obtained from the q-deformed symplectic group with $SU_q(2)$ as structure group, and in [LvS] using 'theta deformations', the structure group being undeformed $SU(2)$. We also mention the coalgebra Galois extension based on quantum coisotropic subgroups given in [BCT, BCDT].

A summary of this paper is as follows. We denote by $\mathcal{H} = \mathcal{O}(SU_q(2))$ the Hopf *-algebra underlying $SU_q(2)$, and by $H = C(SU_q(2))$ the C*-algebraic completion of $\mathcal{H}$. In Section 2 we give a C*-algebra B corresponding to a noncommutative 4-sphere. By construction B is a fibre product $B_0 \times_H B_1$, and can be viewed as a glueing of two noncommutative 3-balls B_0, B_1 over their common boundary H. In fact B is isomorphic to the C*-algebra $C(S^4_q)$ constructed in [DLM] (see also [Pfl]) as a suspension of H. When the parameter q is taken to be 1, then B is commutative and isomorphic to $C(S^4)$, the C*-algebra of continuous functions on the ordinary 4-sphere S^4. The K-groups of B coincide with those of S^4, namely $K_0(B) \cong \mathbb{Z}^2$, $K_1(B) = 0$. One generator of $K_0(B)$ is given by [1], the equivalence class of the unit. The main result of this paper is to give an explicit construction of a family of projections $\{p_n\}_{n\in\mathbb{Z}} \subseteq M_4(B)$, which first of all correspond to powers of a second generator of $K_0(B)$, and secondly are noncommutative analogues of the projections describing classical instanton bundles (1.2). This is achieved in Section 3, as follows. The 6-term cyclic Mayer-Vietoris sequence for K-theory of a fibre product gives an injective map $\partial : K_1(H) \to K_0(B)$, such that $[1] \notin \mathrm{im}(\partial)$. An explicit description of ∂ was given by Bass [Ba] and Milnor [Mi]. Using this, given a generator of $K_1(H)$ (which is known by [MNW90] to be isomorphic to $\mathbb{Z}$), we can construct a second generator of $K_0(B)$. We show, by pairing with an appropriate odd Fredholm module, that $K_1(H)$ is generated by the unitary

$$U = \begin{bmatrix} \alpha & -q\beta^* \\ \beta & \alpha^* \end{bmatrix} \in M_2(H)$$

The projections p_n are related to U^n via $\partial[U^n] = [p_n] - [I_2]$. We give an explicit formula. For each $n \in \mathbb{Z}$, the finitely-generated projective left B-module $B^4 p_n$ is a noncommutative analogue of the classical instanton vector bundle (1.2) with winding number $-n$. Finally we show that the Chern character pairing of $[p_n] \in K_0(B)$ with a suitable even Fredholm module over B is also equal to $-n$.

Similar constructions of projections were given in [CL01] and [DLM] , the former for the quantum 4-sphere $C(S^4_\theta)$. Both correspond to the case of winding number -1. The novelty of our construction is that it provides instanton bundles of arbitrary charges.

Our use of the phrase “noncommutative instanton” to describe certain projections in matrix algebras over noncommutative 4-spheres follows the lead of [CL01]. To fully justify its useage, one should next find a noncommutative analogue of the Yang-Mills action with critical points given by connections with (anti) selfdual curvature. This is a topic for further research.

2 Noncommutative 4-sphere

Throughout this paper we work over the complex numbers as ground field. We denote the coproduct of a Hopf algebra $\mathcal{H}$ using the Sweedler notation $\Delta(h) = \sum h_{(1)} \otimes h_{(2)}$. Here and elsewhere “$\otimes$” denotes the algebraic tensor product (over $\mathbb{C}$). We say that a Hopf algebra $\mathcal{H}$ is a Hopf *-algebra if there exists an antilinear involution $* : \mathcal{H} \to \mathcal{H}$ such that

$$(* \otimes *) \circ \Delta = \Delta \circ *, \quad * \circ S \circ * \circ S = \mathrm{id}, \quad \varepsilon(h^*) = \overline{\varepsilon(h)} \quad \forall\, h \in \mathcal{H}$$

2.1 Quantum $SU(2)$

Recall that $\mathcal{H} = \mathcal{O}(SU_q(2))$, the Hopf $*$-algebra underlying Woronowicz's quantum $SU(2)$ [Wo87], is the unital *-algebra generated by α, β, with relations:

$$\alpha\beta = q\beta\alpha\,,\ \alpha\beta^* = q\beta^*\alpha\,,\ \beta\beta^* = \beta^*\beta\,,\ \alpha^*\alpha + \beta\beta^* = 1\,,\ \alpha\alpha^* + q^2\beta\beta^* = 1\,. \tag{2.3}$$

$\mathcal{H}$ becomes a Hopf *-algebra when equipped with the coproduct, counit and antipode

$$\begin{aligned}&\Delta(\alpha) = \alpha \otimes \alpha - q\,\beta^* \otimes \beta, \quad \Delta(\beta) = \beta \otimes \alpha + \alpha^* \otimes \beta \\ &\varepsilon(\alpha) = 1,\ \varepsilon(\beta) = 0,\ S(\alpha) = \alpha^*,\ S(\beta) = -q\beta,\ S(\beta^*) = -q^{-1}\beta^*,\ S(\alpha^*) = \alpha\end{aligned} \tag{2.4}$$

Let $H = C(SU_q(2))$ be the C*-algebraic completion of $\mathcal{O}(SU_q(2))$, namely the universal C*-algebra generated by indeterminates α, β satisfying the relations (2.3). For $q = 1$, $C(SU_q(2))$ is isomorphic to the commutative unital C*-algebra of continuous functions on the 3-sphere.

2.2 A noncommutative 4-sphere

We now define a C*-algebra B corresponding to the "base space" over which our noncommutative instanton bundles are defined.

Let A be a (unital) C*-algebra, and X a compact Hausdorff space.

Definition 2.1. *Denote by $C(X, A)$ the C*-algebra of continuous functions $f : X \to A$, with pointwise addition and multiplication, and *-structure and norm given by*

$$f^*(x) := f(x)^*, \quad ||f|| := \sup_{x\in X} \{\ ||f(x)||\ \}$$

Definition 2.2. *For $\mathcal{H}$ a *-algebra over $\mathbb{C}$, denote by $C(X, A) \otimes \mathcal{H}$ the algebraic tensor product (over $\mathbb{C}$). This is naturally a *-algebra. For $F \in C(X, A) \otimes \mathcal{H}$, with $F = \sum f^i \otimes h^i$ (finite sum), then for $x \in X$ we denote $F(x) := \sum f^i(x) \otimes h^i \in A \otimes \mathcal{H}$.*

From now on we take $\mathcal{H} = \mathcal{O}(SU_q(2))$, $H = C(SU_q(2))$.

Definition 2.3. *We define C*-algebras*

$$\begin{aligned}&B_0 = \{\ f \in C([0, \tfrac{1}{2}], H) \mid f(0) \in \mathbb{C}1_H\ \}, \quad B_1 = \{\ f \in C([\tfrac{1}{2}, 1], H) \mid f(1) \in \mathbb{C}1_H\ \} \\ &B = \{\ f \in C([0, 1], H) \mid f(0), f(1) \in \mathbb{C}1_H\ \}\end{aligned} \tag{2.5}$$

Let $r_i : B \to B_i$ ($i = 0, 1$) be the obvious restriction maps, and $\pi_i : B_i \to B_{01} := H$ be the projection maps defined by $f \mapsto f(\frac{1}{2})$.

It is immediate that:

Lemma 2.4. *B is a C*-algebra fiber product $B_0 \times_{B_{01}} B_1$, as encoded in the pullback diagram:*

$$\begin{tikzcd} B \arrow[r, "r_1"] \arrow[d, "r_0"'] & B_1 \arrow[d, "\pi_1"'] \\ B_0 \arrow[r, "\pi_0"] & B_{01} \end{tikzcd} \tag{2.6}$$

We note that B is isomorphic to the C*-algebra $C(S^4_q)$ found in [DLM] (see also [Pfl]).

2.3 K-theory of the noncommutative 4-sphere

Applying the Mayer-Vietoris theorem for K-theory of (σ-unital, nuclear) C*-algebras ([Bla], Theorem 21.5.1) to the fiber product $B = B_0 \times_H B_1$ gives the exact sequence:

$$\begin{tikzcd}
K_0(B) \arrow[r, "(r_{0*}, r_{1*})"] & K_0(B_0) \oplus K_0(B_1) \arrow[r, "\pi_{1*} - \pi_{0*}"] & K_0(H) \arrow[d, "\partial"] \\
K_1(H) \arrow[u, "\partial"] & K_1(B_0) \oplus K_1(B_1) \arrow[l, "\pi_{1*} - \pi_{0*}"'] & K_1(B) \arrow[l, "(r_{0*}, r_{1*})"']
\end{tikzcd} \tag{2.7}$$

Lemma 2.5. $K_j(B_i) \cong K_j(\mathbb{C}) = \left\{ \begin{array}{ll} \mathbb{Z} & : j = 0 \\ 0 & : j = 1 \end{array} \right. , \quad i = 0, 1$

Proof. Define $\Sigma_i = \{\ F \in B_i \mid F(i) = 0\ \}$. The sequences $0 \to \Sigma_i \to B_i \to \mathbb{C} \to 0$ are split exact, with splittings $s_i : \mathbb{C} \to B_i$ defined by $(s_i(z))(t) = z$, for all $z \in \mathbb{C}$, $0 \leq t \leq 1$. So $K_j(B_i) \cong K_j(\Sigma_i) \oplus K_j(\mathbb{C})$. Since the Σ_i are contractible their K-groups vanish. So we have isomorphisms $K_j(B_i) \to K_j(\mathbb{C})$ induced by the maps $F \mapsto F(i)$, $i = 0, 1$. In particular $K_0(B_i) \cong \mathbb{Z}$, with generator in each case given by the constant function $\mathbf{1}(t) = 1$ for all t. □

.

Theorem 2.6. $K_0(B) \cong \mathbb{Z}^2$, $K_1(B) = 0$.

Proof. Using Lemma 2.5, the cyclic sequence (2.7) reduces to the exact sequence

$$0 \to K_1(H) \to K_0(B) \to^{(r_{0*}, r_{1*})} K_0(B_0) \oplus K_0(B_1) \to^{\pi_{1*} - \pi_{0*}} K_0(H) \to K_1(B) \to 0$$

Since $K_i(H) \cong \mathbb{Z}$, generated by [1] and $[U]$ (defined in Lemma 2.7) for $i = 0, 1$ respectively, it follows that the map $K_0(B_0) \oplus K_0(B_1) \to K_0(H)$ is surjective, and further $K_0(B) \cong \mathbb{Z}^2$, with one generator given by [1], where 1 is the constant function $1(t) = 1$. Giving an explicit description of a second generator of $K_0(B)$ is the subject of Section 3. □

2.4 K-theory of quantum $SU(2)$

As shown in [MNW90], both K-groups of the C*-algebra $H = C(SU_q(2))$ are isomorphic to $\mathbb{Z}$. The generator of $K_0(H)$ is given by [1]. A result we need in the sequel is:

Lemma 2.7. *Define* $U = \left[\begin{array}{cc} \alpha & -q\beta^* \\ \beta & \alpha^* \end{array} \right] \in M_2(H)$. *Then* $[U]$ *generates* $K_1(H)$.

Proof. Since $U^* = \begin{bmatrix} \alpha^* & \beta^* \\ -q\beta & \alpha \end{bmatrix}$, that $U^*U = I = UU^*$ follows from (2.3). To show U generates $K_1(H)$ we pair with the odd Fredholm module $\mathbf{z}_1$ given in [MNW90], which is defined as follows. Consider the Hilbert space $\ell^2(\mathbb{N}) \otimes \ell^2(\mathbb{Z})$, with basis $\{e_{m,n}\}$. This carries a faithful representation π of H given by

$$\begin{aligned} \pi(\alpha)e_{m,n} &= \lambda_m e_{m-1,n}, \quad \pi(\alpha^*)e_{m,n} = \lambda_{m+1}e_{m+1,n}, \qquad \lambda_m = (1-q^{2m})^{1/2} \\ \pi(\beta)e_{m,n} &= q^m e_{m,n+1}, \quad \pi(\beta^*)e_{m,n} = q^m e_{m,n-1} \end{aligned}$$

Define $Fe_{m,n} = \mathrm{sign}(n)e_{m,n}$. Then $\mathbf{z}_1 = (\ell^2(\mathbb{N}) \otimes \ell^2(\mathbb{Z}), \pi, F)$.

To show U is nontrivial, we compute the Chern character pairing $< [U], \mathrm{ch}_{\mathrm{odd}}(\mathbf{z}_1) >$. Combining the formulae for the Chern character $\mathrm{ch}_{\mathrm{odd}} : KK^1(H, \mathbb{C}) \to HC^{\mathrm{odd}}(\mathcal{H}^\infty)$ and the pairing $< .,. >$ between K-theory and periodic cyclic cohomology given in [Co94], p224 and p296 gives:

$$< [U], \mathrm{ch}_{\mathrm{odd}}(\mathbf{z}_1) >= \lim_{k\to\infty} (-1)^k \, 2^{-(2k+1)} \, \mathrm{Tr}((U^{-1}-1)[\tilde{F}, U]([\tilde{F}, U^{-1}][\tilde{F}, U])^k) \tag{2.8}$$

where $\tilde{F} = \begin{bmatrix} F & 0 \\ 0 & F \end{bmatrix}$. We have

$$[\tilde{F}, U^{-1}][\tilde{F}, U] = \begin{bmatrix} [F,\alpha] & -q[F,\beta^*] \\ [F,\beta] & [F,\alpha^*] \end{bmatrix} \begin{bmatrix} [F,\alpha^*] & [F,\beta^*] \\ -q[F,\beta] & [F,\alpha] \end{bmatrix} = \begin{bmatrix} -q^2 f^* f & 0 \\ 0 & -ff^* \end{bmatrix}$$

for $f = [F, \beta]$, $f^* = [F, \beta]^* = -[F, \beta^*]$. Hence

$$(U^{-1}-1)[\tilde{F}, U]([\tilde{F}, U^{-1}][\tilde{F}, U])^k = (-1)^k \begin{bmatrix} q^{2k+2}\,\beta^* f(f^*f)^k & (1-\alpha)f^*(ff^*)^k \\ q^{2k+1}(1-\alpha^*)f(f^*f)^k & -\beta f^*(ff^*)^k \end{bmatrix}$$

Using the formulae

$$\begin{aligned} q^{2k+2}\,\beta^* f(f^*f)^k e_{m,n} &= \begin{cases} 2^{2k+1}\, q^{2km+2k+2m+2}\, e_{m,-1} & : n = -1 \\ 0 & : n \neq -1 \end{cases}, \\ -\beta f^*(ff^*)^k e_{m,n} &= \begin{cases} -2^{2k+1} q^{2km+2m}\, e_{m,0} & : n = 0 \\ 0 & : n \neq 0 \end{cases} \end{aligned}$$

it follows that

$$\mathrm{Tr}((U^{-1}-1)[\tilde{F}, U]([\tilde{F}, U^{-1}][\tilde{F}, U])^k) = \sum_{m=0}^{\infty} (-1)^k 2^{2k+1} q^{2km+2m}[q^{2k+2}-1] = (-1)^{k+1}2^{2k+1}$$

Hence from (2.8) we have $< [U], \mathrm{ch}_{\mathrm{odd}}(\mathbf{z}_1) >= -1$. By Connes' index theorem [Co94] p296, we have $< x, \mathrm{ch}_{\mathrm{odd}}(\mathbf{z}_1) > \in \mathbb{Z}$ for any $x \in K_1(H)$, so $[U]$ must generate $K_1(H) \cong \mathbb{Z}$.

In [MNW90] it was shown that $K_1(H)$ is generated by $[V]$, where $V = \beta e(0) + (1 - e(0)) \in H$ is a unitary defined in terms of the spectral projection $e(0)$ of $\beta^*\beta$ onto $\{0\}$. Concretely, $e(0)e_{m,n} = \delta_{m,0}e_{m,n}$, hence $Ve_{m,n} = \begin{cases} e_{0,n+1} & : m = 0 \\ e_{m,n} & : m \neq 0 \end{cases}$. It follows from (2.8) that $< [V], \mathrm{ch}_{\mathrm{odd}}(\mathbf{z}_1) >= +1$. Hence $[U] = -[V]$ as elements of $K_1(H)$. □

We thank G. Landi for pointing out to us that the same result was obtained in [DLSVvS2], by computing the index pairing of U with the 3-summable spectral triple found in [DLSVvS1].

3 Noncommutative instantons of arbitrary charge

3.1 The Mayer-Vietoris map on K-theory

Following Bass [Ba] and Milnor [Mi], we give an explicit description of the boundary map $\partial : K_1(B_{01}) \to K_0(B)$ in the Mayer-Vietoris sequence (2.7) associated to a general fiber product of algebras of the form (2.6). This construction defines a map on algebraic K-theory, but it equally well applies to the C*-algebra setting (see e.g. [GH04]). Throughout this Section we assume that the maps $\pi_i : B_i \to B_{01}$ are surjective.

Given a (unital) ring homomorphism $f : R \to S$, and a left R-module E, define f_*E to be the left S-module $S \otimes_R E$. Then there is a canonical R-linear map $f_* : E \to f_*E$ defined by $f_*(e) = 1 \otimes_R e$. It is straightforward to show that, if E is free over R, with basis $\{\hat{x}_\alpha\}$, then f_*E is free over S, with basis $\{x_\alpha := f_*(\hat{x}_\alpha) = 1 \otimes_R \hat{x}_\alpha\}$.

Definition 3.1. *[Mi] Given modules E_0, E_1 over B_0, B_1, then the abelian group $E_0 \times E_1$ is a left B-module via $b \cdot (\xi_0, \xi_1) = (r_0(b)\xi_0, r_1(b)\xi_1)$. Suppose we are given an isomorphism $h : \pi_{0*}E_0 \cong \pi_{1*}E_1$. Define $E = M(E_0, E_1, h)$ to be the submodule of $E_0 \times E_1$ consisting of all pairs (ξ_0, ξ_1) such that $h \circ \pi_{0*}(\xi_0) = \pi_{1*}(\xi_1)$.*

The following is a special case of [Mi], Theorems 2.1-2.3, although the formula for the idempotent p is not given there.

Theorem 3.2. *Suppose that E_0 and E_1 are free over B_0, B_1 and both of rank N. Then E is finitely generated projective over B. Explicitly, $E \cong B^{2N}p$, where $p \in M_{2N}(B)$ is given by*

$$p = \begin{bmatrix} (1, c(2-dc)d) & (0, c(2-dc)(1-dc)) \\ (0, (1-dc)d) & (0, (1-dc)^2) \end{bmatrix} \tag{3.9}$$

for elements c, $d \in M_N(B_1)$ defined below.

Proof. For clarity we give a self-contained proof. Suppose that $E_0 \cong B_0^N$, $E_1 \cong B_1^N$, with bases $\{\hat{x}_\alpha\}_{1\le\alpha\le N}$, $\{\hat{y}_\beta\}_{1\le\beta\le N}$ respectively, and corresponding bases $\{x_\alpha := \pi_{0*}(\hat{x}_\alpha)\}$, $\{y_\beta := \pi_{1*}(\hat{y}_\beta)\}$ for $\pi_{0*}E_0$, $\pi_{1*}E_1$, which are both isomorphic to B_{01}^N. The isomorphism $h : \pi_{0*}E_0 \cong \pi_{1*}E_1$ is defined by $h(x_\alpha) = \sum a_{\alpha\beta} y_\beta$, for some invertible $a = (a_{\alpha\beta}) \in M_N(B_{01})$. Fix a lifting c of a under π_1, i.e. a matrix $c = (c_{\alpha\beta}) \in M_N(B_1)$, such that $\pi_1(c_{\alpha\beta}) = a_{\alpha\beta}$ for all α, β.

Case 1. Suppose that c is invertible. For each α, define $\hat{y}'_\alpha := \sum c_{\alpha\beta}\hat{y}_\beta$. Since c is invertible, then $\{\hat{y}'_\alpha\}$ is a basis for E_1. Since $h \circ \pi_{0*}(\hat{x}_\alpha) = h(x_\alpha) = \sum a_{\alpha\beta} y_\beta = \pi_{1*}(\hat{y}'_\alpha)$, it follows that $\hat{z}_\alpha := (\hat{x}_\alpha, \hat{y}'_\alpha) \in E = M(E_0, E_1, h)$ for all α. It is then straightforward to check that $\{\hat{z}_\alpha\}_{1\le\alpha\le N}$ span E and are linearly independent, i.e. E is free, $E \cong B^N$ with basis $\{\hat{z}_\alpha\}$.

Case 2. Suppose that c is not invertible. Fix a lifting $d = (d_{\alpha\beta}) \in M_N(B_1)$ of a^{-1}. Define free modules $F_0 \cong B_0^N$, $F_1 \cong B_1^N$, with bases $\{\hat{u}_\alpha\}$, $\{\hat{v}_\beta\}$. As before $\pi_{0*}F_0$, $\pi_{1*}F_1$ are both isomorphic to B_{01}^N, with bases $\{u_\alpha\}$, $\{v_\beta\}$. Define $g : \pi_{0*}F_0 \cong \pi_{1*}F_1$ by $g(u_\alpha) = \sum a^{-1}_{\alpha\beta} v_\beta$. Then $h \oplus g$ defines an isomorphism $\pi_{0*}(E_0 \oplus F_0) \cong \pi_{1*}(E_1 \oplus F_1)$. Using Whitehead's lemma we write

$$\begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix} = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -a^{-1} & 1 \end{bmatrix} \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \in M_{2N}(B_{01})$$

which lifts to

$$\mathcal{C} = \begin{bmatrix} \mathcal{C}^{11} & \mathcal{C}^{12} \\ \mathcal{C}^{21} & \mathcal{C}^{22} \end{bmatrix} = \begin{bmatrix} 1 & c \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -d & 1 \end{bmatrix} \begin{bmatrix} 1 & c \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} (2-cd)c & cd-1 \\ 1-dc & d \end{bmatrix} \in M_{2N}(B_1)$$

In our notation, each $\mathcal{C}^{ij} = (\mathcal{C}^{ij}_{\alpha\beta}) \in M_N(B_1)$. Note that $\mathcal{C}$ is invertible, with inverse

$$\mathcal{D} = \begin{bmatrix} \mathcal{D}^{11} & \mathcal{D}^{12} \\ \mathcal{D}^{21} & \mathcal{D}^{22} \end{bmatrix} = \begin{bmatrix} d & 1-dc \\ cd-1 & (2-cd)c \end{bmatrix} \in M_{2N}(B_1)$$

hence Case 1 applies. We have bases $\{\hat{x}_\alpha, \hat{u}_\beta\}_{1\le\alpha,\beta\le N}$, $\{\hat{y}_\alpha, \hat{v}_\beta\}_{1\le\alpha,\beta\le N}$ for $E_0 \oplus F_0$, $E_1 \oplus F_1$ respectively. Define

$$\hat{y}'_\alpha := \sum \mathcal{C}^{11}_{\alpha\alpha'}\ \hat{y}_{\alpha'} + \sum \mathcal{C}^{12}_{\alpha\beta'}\ \hat{v}_{\beta'}\ , \quad \hat{v}'_\beta := \sum\ \mathcal{C}^{21}_{\beta\alpha'}\ \hat{y}_{\alpha'} + \sum \mathcal{C}^{22}_{\beta\beta'}\ \hat{v}_{\beta'}$$

By the above, $M(E_0 \oplus F_0, E_1 \oplus F_1, h \oplus g)$ is free over B of rank $2N$, with basis $\{(\hat{x}_\alpha, \hat{y}'_\alpha), (\hat{u}_\beta, \hat{v}'_\beta)\}_{1\le\alpha,\beta\le N}$, and further

$$M(E_0, E_1, h) \oplus M(F_0, F_1, g) \cong M(E_0 \oplus F_0, E_1 \oplus F_1, h \oplus g)$$

via $((e_0, e_1), (f_0, f_1)) \mapsto ((e_0, f_0), (e_1, f_1))$. So $E = M(E_0, E_1, h)$ is a direct summand of a free module and hence projective.

So we have a projection and inclusion

$$M(E_0 \oplus F_0, E_1 \oplus F_1, h \oplus g) \to^{\pi} M(E_0, E_1, h) \hookrightarrow^{i} M(E_0 \oplus F_0, E_1 \oplus F_1, h \oplus g)$$

with $(i \circ \pi)^2 = i \circ \pi$. We need to find the matrix $p \in M_{2N}(B)$ corresponding to $i \circ \pi$ with regard to the basis $\{(\hat{x}_\alpha, \hat{y}'_\alpha), (\hat{u}_\beta, \hat{v}'_\beta)\}$. We have

$$\begin{aligned} i \circ \pi(\hat{x}_\alpha, \hat{y}'_\alpha) &= i \circ \pi(\hat{x}_\alpha, \sum \mathcal{C}^{11}_{\alpha\alpha'}\ \hat{y}_{\alpha'} + \sum \mathcal{C}^{12}_{\alpha\beta'}\ \hat{v}_{\beta'}) = (\hat{x}_\alpha, \sum \mathcal{C}^{11}_{\alpha\alpha'}\ \hat{y}_{\alpha'}) \\ i \circ \pi(\hat{u}_\beta, \hat{v}'_\beta) &= i \circ \pi(\hat{u}_\beta, \sum\ \mathcal{C}^{21}_{\beta\alpha'}\ \hat{y}_{\alpha'} + \sum \mathcal{C}^{22}_{\beta\beta'}\ \hat{v}_{\beta'}) = (0, \sum\ \mathcal{C}^{21}_{\beta\alpha'}\ \hat{y}_{\alpha'}) \end{aligned} \tag{3.10}$$

We show this corresponds to the matrix

$$p = \begin{bmatrix} P^{11} & P^{12} \\ P^{21} & P^{22} \end{bmatrix} = \begin{bmatrix} (1, c(2-dc)d) & (0, c(2-dc)(1-dc)) \\ (0, (1-dc)d) & (0, (1-dc)^2) \end{bmatrix} = \begin{bmatrix} (1, Q^{11}) & (0, Q^{12}) \\ (0, Q^{21}) & (0, Q^{22}) \end{bmatrix}$$

We have

$$\begin{aligned} p(\hat{x}_\alpha, \hat{y}'_\alpha) &= \sum\ P^{11}_{\alpha\alpha'}(\hat{x}_{\alpha'}, \hat{y}'_{\alpha'}) + \sum\ P^{12}_{\alpha\beta}(\hat{u}_\beta, \hat{v}'_\beta) = (\hat{x}_\alpha, \sum Q^{11}_{\alpha\alpha'}\ \hat{y}'_{\alpha'}) + (0, \sum\ Q^{12}_{\alpha\beta}\ \hat{v}'_\beta) \\ &= (\hat{x}_\alpha, \sum\ Q^{11}_{\alpha\alpha'}\ \mathcal{C}^{11}_{\alpha'\alpha''}\ \hat{y}_{\alpha''} + \sum\ Q^{11}_{\alpha\alpha'}\ \mathcal{C}^{12}_{\alpha'\beta'}\ \hat{v}_{\beta'} + \sum\ Q^{12}_{\alpha\beta}\ \mathcal{C}^{21}_{\beta\alpha'}\ \hat{y}_{\alpha'} + \sum\ Q^{12}_{\alpha\beta}\ \mathcal{C}^{22}_{\beta\beta'} \hat{v}_{\beta'}) \\ &= (\hat{x}_\alpha, \sum\ (Q^{11}\mathcal{C}^{11} + Q^{12}\mathcal{C}^{21})_{\alpha\alpha'}\ \hat{y}_{\alpha'} + \sum\ (Q^{11}\mathcal{C}^{12} + Q^{12}\mathcal{C}^{22})_{\alpha\beta'}\ \hat{v}_{\beta'}) \end{aligned}$$

Now,

$$\begin{aligned} Q^{11}\mathcal{C}^{11} + Q^{12}\mathcal{C}^{21} &= c(2-dc)^2 dc + c(2-dc)(1-dc)^2 = c(2-dc) = \mathcal{C}^{11}, \\ Q^{11}\mathcal{C}^{12} + Q^{12}\mathcal{C}^{22} &= cd(2-cd)(cd-1) + cd(2-cd)(1-cd) = 0 \end{aligned}$$

Hence $p(\hat{x}_\alpha, \hat{y}'_\alpha) = (\hat{x}_\alpha, \sum \mathcal{C}^{11}_{\alpha\alpha'}\, \hat{y}_{\alpha'})$, which agrees with (3.10). In exactly the same way it can be checked that $p(\hat{u}_\beta, \hat{v}'_\beta) = (0, \sum \mathcal{C}^{21}_{\beta\alpha'}\, \hat{y}_{\alpha'})$. Hence p is the matrix for $i \circ \pi$. $\square$

The Mayer-Vietoris boundary map $\partial : K_1(B_{01}) \to K_0(B)$ can now be described explicitly [Ba, Mi]. Given an invertible matrix $V \in M_N(B_{01})$ representing an element of $K_0(B_{01})$, take $E_i := B_i^N$ for $i = 0, 1$, then V defines an isomorphism $\pi_{0*}E_0 \cong \pi_{1*}E_1$, both of which are isomorphic to B_{01}^N. Then $E = M(E_0, E_1, V)$ is finitely generated projective, $E \cong B^{2N}p$, where $p \in M_{2N}(B)$ is the corresponding projection defined by (3.9). Then $\partial[V] = [p] - [I_N]$, where I_N is the rank N identity matrix.

3.2 Instanton projections of arbitrary charge

The unitary $U \in M_2(H)$ defined in Lemma 2.7 generates $K_1(H) \cong \mathbb{Z}$. In particular, for each $n \in \mathbb{Z}$, $[U^n] = n[U] \in K_1(H)$.

Definition 3.3. *For each $n \in \mathbb{Z}$, let $p_n \in M_4(B)$ be the projection corresponding to U^n defined by (3.9). We will refer to p_n as the "instanton projection of charge $-n$".*

Theorem 3.4. *For each $n \in \mathbb{Z}$, the projection $p_n \in M_4(B)$ is explicitly given by*

$$\begin{aligned} n \geq 0: \quad p_n &= \begin{bmatrix} c^n(2 - d^nc^n)d^n & c^n(2 - d^nc^n)(1 - d^nc^n) \\ (1 - d^nc^n)d^n & (1 - d^nc^n)^2 \end{bmatrix} \\ n \leq 0: \quad p_{-n} &= \begin{bmatrix} d^n(2 - c^nd^n)c^n & d^n(2 - c^nd^n)(1 - c^nd^n) \\ (1 - c^nd^n)c^n & (1 - c^nd^n)^2 \end{bmatrix} \end{aligned} \tag{3.11}$$

Proof. Using Theorem 3.2, first U, U^{-1} lift to $c = \begin{bmatrix} \bar{\alpha} & -q\bar{\beta}^* \\ \bar{\beta} & \bar{\alpha}^* \end{bmatrix}$, $d = \begin{bmatrix} \bar{\alpha}^* & \bar{\beta}^* \\ -q\bar{\beta} & \bar{\alpha} \end{bmatrix} \in M_2(B)$, where $\overline{\alpha}$, $\overline{\beta} \in B$ are defined by:

$$\overline{\alpha}(t) = \begin{cases} (2t-1) + 2(1-t)\alpha & : \frac{1}{2} \leq t \leq 1 \\ \alpha & : 0 \leq t \leq \frac{1}{2} \end{cases}, \quad \overline{\beta}(t) = \begin{cases} 2(1-t)\beta & : \frac{1}{2} \leq t \leq 1 \\ \beta & : 0 \leq t \leq \frac{1}{2} \end{cases} \tag{3.12}$$

So for $n \geq 0$, U^n lifts to c^n, and U^{-n} lifts to d^n. Note that for $0 \leq t \leq \frac{1}{2}$ we have

$$[c(2 - dc)d](t) = I_2, \quad [c(2 - dc)(1 - dc)](t) = 0 = [(1 - dc)d](t) = [(1 - dc)^2](t)$$

The result then follows from (3.9). $\square$

Dual to the Mayer-Vietoris six-term cyclic sequence on K-theory (2.7) is a cyclic sequence on K-homology ([Bla] Theorem 21.5.1):

$$\begin{tikzcd} K^0(B) \arrow[d, "\delta"'] & K^0(B_0) \oplus K^0(B_1) \arrow[l, "r_0^* + r_1^*"'] & K^0(H) \arrow[l, "(-\pi_0^*, \pi_1^*)"'] \\ K^1(H) \arrow[r, "(-\pi_0^*, \pi_1^*)"] & K^1(B_0) \oplus K^1(B_1) \arrow[r, "r_0^* + r_1^*"] & K^1(B) \arrow[u, "\delta"'] \end{tikzcd} \tag{3.13}$$

where $K^j(D)$ denotes the Kasparov K-homology group $KK^j(D,\mathbb{C})$. By Rosenberg and Schochet's universal coefficient theorem, if the K-groups $K_j(D)$ are free, then $K^j(D) \cong K_j(D)$ (as abelian groups). So for our fibre product $B = B_0 \times_H B_1$, we have

$$K^j(H) = \begin{cases} \mathbb{Z} & : j=0 \\ \mathbb{Z} & : j=1 \end{cases}, \quad K^j(B_i) = \begin{cases} \mathbb{Z} & : j=0 \\ 0 & : j=1 \end{cases}, \quad K^j(B) = \begin{cases} \mathbb{Z}^2 & : j=0 \\ 0 & : j=1 \end{cases},$$

(the Fredholm modules generating $K^j(H)$ were given in [MNW90]). Hence $\delta : K^0(B) \to K^1(H)$ is surjective. By Lemma 2.7, we have that $K_1(H)$ is generated by $[U]$, and $K^1(H)$ generated by the equivalence class of the Fredholm module $\mathbf{z}_1$. Now, $[U] \in KK^1(\mathbb{C}, H)$, $\mathbf{z}_1 \in KK^1(H,\mathbb{C})$, and the pairing $< [U], \mathrm{ch}_{\mathrm{odd}}(\mathbf{z}_1) >$ is given by the Kasparov product $[U]\hat{\otimes}_H \mathbf{z}_1 \in KK^0(\mathbb{C},\mathbb{C}) \cong \mathbb{Z}$.

It follows from [Bla, Scho] that the Mayer-Vietoris boundary maps ∂ and δ are dual, in the sense that, for any $\mathbf{w}_0 \in K^0(B)$ such that $\delta(\mathbf{w}_0) = \mathbf{z}_1 \in K^1(H)$, then

$$< \partial[x], \mathrm{ch}_{\mathrm{even}}(\mathbf{w}_0) >=< [x], \mathrm{ch}_{\mathrm{odd}}(\mathbf{z}_1) > \quad \forall\ [x] \in K_1(H)$$

Hence

$$< p_n, \mathrm{ch}_{\mathrm{even}}(\mathbf{w}_0) >=< \partial[U^n], \mathrm{ch}_{\mathrm{even}}(\mathbf{w}_0) >=< [U^n], \mathrm{ch}_{\mathrm{odd}}(\mathbf{z}_1) >= n < [U], \mathrm{ch}_{\mathrm{odd}}(\mathbf{z}_1) >= -n$$

thus justifying the terminology "instanton projection of charge $-n$".

Acknowledgments. We thank G. Landi, R. Nest, M. Wodzicki and S. L. Woronowicz for their very useful comments.

TH: I thank Katedra Metod Matematycznych Fizyki, Uniwersytet Warszawski for their support via EU Transfer of Knowledge contract MKTD-CT-2004-509794. I am also very grateful to Instytut Matematyczny, Polska Akademia Nauk, and to the Isaac Newton Institute, Cambridge, for their support and hospitality.

References

[At] M. F. Atiyah, *The geometry of Yang-Mills fields*, Fermi Lectures, Scuola Normale Superiore Pisa, Pisa (1979).

[Ba] H. Bass, *Algebraic K-theory*, W.A. Benjamin, Inc., New York-Amsterdam (1968).

[Bla] B. Blackadar, *K-theory for operator algebras*, MSRI publications **5**, Cambridge University Press (1998).

[BCT] F. Bonechi, N. Ciccoli, M. Tarlini, *Noncommutative instantons on the 4-sphere from quantum groups*, Commun. Math. Phys. **226** (2002) 419-432.

[BCDT] F. Bonechi, N. Ciccoli, L. Dąbrowski, M. Tarlini, *Bijectivity of the canonical map for the noncommutative instanton bundle*, J. Geom. Phys. **51** (2004) 419-432.

[Co94] A. Connes, *Noncommutative geometry*, Academic Press, San Diego (1994).

[CL01] A. Connes, G. Landi, *Noncommutative manifolds, the instanton algebra and isospectral deformations*, Commun. Math. Phys. **221**, 141-159 (2001).

[DHH] L. Dąbrowski, T. Hadfield, P. M. Hajac, *Simplicial constructions of principal extensions*, in preparation (2006).

[DLM] L. Dąbrowski, G. Landi, T. Masuda, *Instantons on the Quantum 4-Spheres* S^4_q, Commun. Math. Phys. **221** (2001) 161-168.

[DLSVvS1] L. Dąbrowski, G. Landi, A. Sitarz, J. C. Vrilly, W. van Suijlekom, *The Dirac operator on* $\mathrm{SU}_q(2)$. Commun. Math. Phys. **259**, 729-759 (2005).

[DLSVvS2] L. Dąbrowski, G. Landi, A. Sitarz, J. C. Vrilly, W. van Suijlekom, *The local index formula for* $\mathrm{SU}_q(2)$. K-Theory **35**, no. 3-4, 375-394 (2005).

[GH04] E. Guentner, N. Higson, *Group C*-algebras and K-theory.* Noncommutative geometry, 137-251, Lecture Notes in Mathematics **1831**, Springer, Berlin (2004).

[KS97] A. Klimyk, K. Schmüdgen, *Quantum groups and their representations*, Springer (1997).

[LPR] G. Landi, C. Pagani, C. Reina, *A Hopf bundle over a quantum four-sphere from the symplectic group*, Commun. Math. Phys. **263** (2006) 65-88.

[LvS] G. Landi, W. van Suijlekom, *Principal fibrations from noncommutative spheres*, Commun. Math. Phys. **260** (2005) 203-225.

[Mi] J. Milnor, *Introduction to algebraic K-theory.* Annals of Mathematics Studies, **72**. Princeton University Press, Princeton, N.J. University of Tokyo Press, Tokyo, 1971.

[MNW90] T. Masuda, Y. Nakagami, J. Watanabe, *Noncommutative differential geometry on the quantum* $SU(2)$*, I,* K-theory **4**, no. 2, 157-180 (1990).

[Mat] K. Matsumoto, *Noncommutative three-dimensional spheres.* Japan. J. Math. (N.S.) **17** (1991), no. 2, 333–356.

[Pfl] M.J. Pflaum, *Quantum Groups on Fibre Bundles*, Commun. Math. Phys. **166** (1994) 279-316.

[Scho] C. Schochet, *Topological methods for C*-algebras, III : Axiomatic homology.* Pacific J. Math. **114** no. 2, 399-445 (1984).

[Wo87] S.L. Woronowicz, *Twisted* $SU(2)$ *group : an example of a noncommutative differential calculus,* Publ. R.I.M.S. (Kyoto University) **23**, 117-181 (1987).